Original Article

Evaluation of the effects of a low dose of *Asiasari radix* on stem cell morphology and proliferation

Jun-Beom Park¹, Ji-Eun Lee¹, Seong-Ho Jin¹, Youngkyung Ko¹, Su-Hyeon Jeong²

¹Department of Periodontics, College of Medicine, The Catholic University of Korea

²Department of Rehabilitation Medicine of Korean Medicine, Chungju Hospital of Korean Medicine, College of Korean Medicine, Semyung University

Objectives: *Asiasari radix* (*A. radix*) is a traditional herb medicine that has been used as an analgesic, antitussive, or anti-allergic remedy. This study was performed to evaluate the effects of low concentration of the *Asiasarum heterotropoides* extract on the morphology and proliferation of the human mesenchymal stem cells derived from periodontal tissue.

Methods: Stem cells derived from gingiva were grown in the presence of *A. radix* at final concentrations that ranged from 0.001 to 0.01 $\mu\text{g/mL}$. The morphology of the cells was viewed under an inverted microscope and the analysis of cell proliferation was performed by using Cell Counting Kit-8 (CCK-8) on days 1 and 3.

Results: The control group showed fibroblast morphology. The shapes of the cells in 0.001 and 0.01 $\mu\text{g/mL}$ of *A. radix* were similar to that of the untreated control group. The cultures growing in the presence of *A. radix* at day 1 showed an increase in the CCK-8 value. The relative values of CCK-8 assays of 0.001 and 0.01 $\mu\text{g/mL}$ of *A. radix* are $130.6\% \pm 1.8\%$ and $129.3\% \pm 1.5\%$, respectively, when the CCK-8 result of the untreated control group at day 1 is considered 100% ($P = 0.051$).

Conclusions: Within the limits of this study, low concentrations of *A. radix* seemed to increase the proliferation of the stem cells that were derived from the gingiva and did not have adverse effects on the morphology of the cells.

Key Words : *Asiasari Radix*, low dose, stem cells, cell morphology, cell proliferation

Introduction

Asiasari Radix (*A. radix*) is a traditional herb that is widely used to treat various diseases in Korea and China¹. *A. radix* is primarily derived from *Asiasarum heterotropoides* or *Asiasarum sieboldii*²⁻³. It has been used as an analgesic, antitussive, or anti-allergic remedy for generations⁴. It has also been used to treat dental diseases that include aphthous stomatitis, toothache, and gingivitis^{1,5}. In Dongeubogam, there

are many prescriptions of main blended *A. radix*, and those were the most frequently used to treat dental diseases that included gingivitis⁶. However, only limited information is currently available regarding the effects of *A. radix* on dental tissue, including mesenchymal stem cells derived from gingiva.

Several studies were performed to evaluate the effects of *A. radix* on cell proliferations with conflicting results^{3-5,7,8}. *A. radix* had no significant

• Received : 26 May 2016 • Revised : 17 June 2016 • Accepted : 24 June 2016
• Correspondence to : Su-Hyeon Jeong
Department of Rehabilitation Medicine of Korean Medicine,
Chungju Hospital of Korean Medicine, College of Korean Medicine, Semyung University
Sangbang 4gil, Chung ju, Chung buk, 27429, Republic of Korea
Tel : +82-43-841-1737, Fax : +82-43-856-1731, E-mail : js365a@hanmail.net

effects on the growth of HeLa cells at a range of concentrations (0.0001 to 1,000 $\mu\text{g/mL}$) at 72 hr⁷⁾. Similarly, neither the proliferation nor the viability of whole spleen cells or human U266B1 multiple myeloma cells were affected with 1000 $\mu\text{g/mL}$ of *A. radix*⁴⁾. However, in a study on a human colorectal carcinoma cell line (HCT-116 cells), increasing concentrations (0-50 $\mu\text{g/mL}$) of ethanol extract of *A. radix* for 48 h were shown to repress the cell proliferation in a dose-dependent manner⁸⁾. Similarly, our previous study showed that the groups treated with 100 and 1,000 $\mu\text{g/mL}$ (tested range of concentrations of 0.1 to 1,000 $\mu\text{g/mL}$) repressed the proliferation³⁾. On the other hand, *A. radix* at a dose of 0.1 $\mu\text{g/mL}$ increased the proliferation of an immortalized human keratinocyte cell line and the primary cultures of human hair dermal papilla cells⁵⁾.

Limited information is currently available regarding the effects of *A. radix* on the oral tissue³⁾. Our groups also evaluated the effects of other traditional herb including Angelicae dahuricae radix and Cimicifugae Rhizoma on mesenchymal stem cells derived from gingiva^{9,10)}. This study was performed to evaluate the effects of the extract of *Asiasarum heterotropoides* at a low concentration on the morphology and proliferation of the human mesenchymal stem cells derived from periodontal tissue.

Methodology

1. Preparation of the materials

The dry roots of *Asiasarum heterotropoides* (400 g) were immersed in distilled water and boiled under reflux for 2 h 30 min. The resulting extract was centrifuged and the supernatant was concentrated to 300 mL by using a rotary evaporator (Eyela NE-1001, Toky Rikakikai Co. Ltd, Tokyo, Japan) under reduced pressure. The concentrates were then freeze-dried in lyophilizer (Labconco, Kansas, USA). Sixty-five gram of solid residue was obtained and

the yield was 16% (w/w).

2. Isolation and culture of the stem cells derived from the gingiva

Healthy gingival tissue samples were obtained from healthy patient during crown lengthening procedures. Crown lengthening procedures are performed to provide adequate room for crown preparation and reestablishment of the biologic width between the crown margin and the bone crest¹¹⁾. This study was reviewed and approved by the Institutional Review Board of Seoul St. Mary's Hospital, College of Medicine, The Catholic University of Korea, Seoul, Republic of Korea (KC11SISI0348), and informed consents were obtained from the patients.

The resected gingival tissues were immediately placed in sterile phosphate-buffered saline (PBS, Welgene, Daegu, Korea) containing penicillin, and streptomycin (Sigma-Aldrich Co., St. Louis, USA) at 4°C. The tissue was de-epithelialized, and digested with collagenase IV (Sigma-Aldrich Co.). Then the cells were incubated at 37°C in a humidified incubator. The non-adherent cells were washed with PBS (Welgene), replaced with fresh medium, and fed every 2-3 days.

3. Evaluation of stem cell morphology

The stem cells were plated at a density of 2.0×10^3 cells/well in 96-well plates. The cells were incubated in Minimum Essential Medium α (α -MEM, Gibco, Grand Island, USA) with 10 mM ascorbic acid 2-phosphate (Sigma-Aldrich Co.), 200 mM L-Glutamine (Sigma-Aldrich Co.), 15% fetal bovine serum (Gibco), 100 U/mL penicillin, and 100 $\mu\text{g/mL}$ streptomycin (Sigma-Aldrich Co.) in the presence of the *A. radix* at final concentrations that ranged from 0 to 0.01 $\mu\text{g/mL}$ (0 (untreated control), 0.001 and 0.01 $\mu\text{g/mL}$), respectively. The morphology of the cells was viewed under an inverted microscope (Leica DM IRM, Leica Microsystems, Wetzlar, Germany) on days 1 and 3.

4. Determination of cell proliferation

The analysis of cell proliferation was performed on days 1 and 3. Cell Counting Kit-8 (CCK-8, Dojindo, Tokyo, Japan) assay was used to identify viable cells. The spectrophotometric absorbance was measured at 450 nm with a microplate reader (BioTek, Winooski, USA). The experiments were done in triplicate.

5. Statistical Analysis

The findings are represented as the means $\pm$ standard deviations of the experiments. A test of normality was performed and Levene's test of homogeneity of variances revealed the *P*-value of

0.040. Non-parametric statistical procedures with Kruskal-Wallis methods were used to test for differences between three groups with commercially available statistical software (SPSS 12 for Windows, SPSS Inc., Chicago, IL, USA).

Results

1. Evaluation of cell morphology

These cells showed colony-forming abilities, plastic adherence, and multilineage differentiation (osteogenic, adipogenic, chondrogenic) potency and expressed CD44, CD73, CD90, and CD105, but did not express CD14, CD45, CD34, and CD19 in flow cytometry. The control group showed fibroblast-like

Fig. 1. Evaluation of cell morphology on day 1.

The control group showed fibroblast morphology on day 1. The shapes of the cells in 0.001 and 0.01 $\mu\text{g/mL}$ of *Asiasari radix* were similar to that of the untreated control group.

A : Control group

B : 0.001 $\mu\text{g/mL}$ of *Asiasari radix*

C : 0.01 $\mu\text{g/mL}$ of *Asiasari radix*

Fig. 2. Cell morphology on day 3.

The shapes of the cells in 0.001 and 0.01 $\mu\text{g/mL}$ of *Asiasari radix* were similar to that of the untreated control group.

A : Control group

B : 0.001 $\mu\text{g/mL}$ of *Asiasari radix*

C : 0.01 $\mu\text{g/mL}$ of *Asiasari radix*

morphology on day 1 (Fig. 1). The shapes of the cells in 0.001 and 0.01 $\mu\text{g/mL}$ of *A. radix* were similar to that of the untreated control group. The morphology of the cells on day 3 is shown in Figure 2. The shapes of the cells in 0.001 and 0.01 $\mu\text{g/mL}$ were similar to that of the untreated control group.

2. Cell proliferation

The results of cell proliferation on days 1 and 3 are shown in Figures 3 and 4, respectively. The cultures that were growing in the presence of *A. radix* on day 1 showed an increase in the CCK-8 value. The relative values of CCK-8 assays of 0.001 and 0.01 $\mu\text{g/mL}$ of *A. radix* are $130.6 \% \pm 1.8 \%$ and $129.3 \% \pm 1.5 \%$, respectively, when the CCK-8 result of the untreated control group on day 1 is

Fig. 3. Cellular viability on day 1.

The cultures that were growing in the presence of *Asiasari radix* on day 1 showed an increase in the CCK-8 value. The relative values of CCK-8 assays of 0.001 and 0.01 $\mu\text{g/mL}$ of *Asiasari radix* are $130.6 \% \pm 1.8 \%$ and $129.3 \% \pm 1.5 \%$, respectively, when the CCK-8 result of the untreated control group on day 1 is considered 100% ($100.0 \% \pm 21.9 \%$) ($P = 0.051$).

considered 100% ($100.0 \% \pm 21.9 \%$) ($P = 0.051$). The results on day 3 are shown in Figure 4.

The cultures that were growing in the presence of 0.001 and 0.01 $\mu\text{g/mL}$ of *A. radix* did not show any statistically significant changes in the CCK-8 values on day 3 ($P = 0.288$). The relative values of 0.001 and 0.01 $\mu\text{g/mL}$ groups are $96.5 \% \pm 2.0 \%$ and $98.6 \% \pm 2.5 \%$, respectively, when the CCK-8 results of the control group on day 3 is considered 100 % ($100.0 \% \pm 3.5 \%$).

Discussion

In this report, we examined the effects of a low concentration of the extract of *Asiasari radix* on the morphology and proliferation of the human mesenchymal stem cells that were derived from periodontal tissue. Low concentrations (0.001 and 0.01 $\mu\text{g/mL}$) of *A. radix* seemed to increase the proliferation of the stem cells derived from the gingiva.

Previous studies evaluating the effects of *A. radix* on cell proliferations showed conflicting results. Some showed no significant changes at the range of concentrations (0.0001 to 1,000 $\mu\text{g/mL}$)^{4,7}, but others showed repressed cell proliferation in a dose-dependent manner⁸. However, in another report, the *A. radix* extract at a dose of 0.1 $\mu\text{g/mL}$ increased the proliferation⁵. The different responses to *A. radix* may be explained in part, by the type of cells, culture condition, or culture period^{3,12}.

Mesenchymal stem cells derived from gingival connective tissue have the capacity to be differentiated into osteoblasts, chondroblasts, and adipocytes¹³, and have immunomodulatory anti-inflammatory and regenerative properties¹⁴⁻¹⁶. They are relatively homogenous and they proliferate fast and they are reported to have easy, relatively noninvasive access¹⁷⁻¹⁸. Moreover, gingiva-derived mesenchymal stem cells were suggested to be superior to bone marrow-derived mesenchymal stem cells in that they do not lose the MSC characteristic at higher

Fig. 4. Cellular viability on day 3.

The cultures that were growing in the presence of 0.001 and 0.01 $\mu\text{g/mL}$ of *Asiasari radix* did not show any statistically significant changes in the CCK-8 assays on day 3 ($P = 0.288$). The relative values of 0.001 and 0.01 $\mu\text{g/mL}$ groups are $96.5 \% \pm 2.0 \%$ and $98.6 \% \pm 2.5 \%$, respectively, when the CCK-8 results of the untreated control group on day 3 is considered 100 % ($100.0 \% \pm 3.5 \%$).

passages, maintain a normal karyotype and exhibit telomerase activity in long-term cultures¹⁸. These stem cells may be very useful in the research field and in the treatment of disease^{3,13}. Stem cells derived from the gingiva have been applied for the treatment of rheumatoid arthritis, colitis, and allergic contact dermatitis^{15,19,20}, bony and cutaneous wound healing^{21,22}. There also is a study demonstrating new bone growth from the transplanted gingiva-derived stem cells in the calvarial and the mandibular rat defect²². Gingiva-derived stem cell spheroids were fabricated using concave microwells and the shape and the viability were maintained during the experimental periods²³. It can be suggested that gingiva-derived stem cells represent an accessible candidate for regenerative therapies²⁴.

Various methods are available for the assessment of cellular proliferation²⁵⁻³⁰. Bradford assay may be used as an indirect measurement of cell proliferation because it measures the content of viable cells by measuring total protein concentration²⁶. Trypan blue assay is a dye exclusion test that is used to determine the number of viable cells because live cells possess

intact cell membranes that exclude the dyes²⁵). The MTT (3-(4,5-Dimethylthiazol-2-yl)-2,5-diphenyltetrazolium bromide) assay may be more sensitive than trypan blue because it assesses the cell proliferation through the determination of mitochondrial dehydrogenase activity²⁷). However, an additional step is needed for the MTT assay; this step requires the water insoluble formazan salt to be solubilized for the assay²⁸). The tetrazolium dyes MTS and XTT form a soluble formazan that can be measured directly without an additional solubilization step²⁹). CCK-8 assay used in this study relies on the ability of mitochondrial dehydrogenases to oxidize WST-8 [2-(2-methoxy-4-nitrophenyl)-3-(4-nitrophenyl)-5-(2,4-disulfophenyl)-2H tetrazolium, monosodium salt] into a formazan product³⁰). In this study, the CCK-8 assay was used because the detection sensitivity of CCK-8 is higher than other assays using MTT, XTT or MTX tetrazolium dyes and tetrazolium dye WST-8 in CCK-8 assay is reported to be less toxic to the tested cells³¹⁻³²). Proliferation may also be assessed with the assay, which is based on the detection of 5'-bromo-2'-deoxy-uridine (BrdU) that is incorporated into the DNA of proliferating cells in place of thymidine³²).

Within the limits of this study, low concentrations of *A. radix* seemed to increase the proliferation of the stem cells that were derived from the gingiva and did not have adverse effects on the morphology of the cells.

Acknowledgement

This paper was supported by the Semyung University Research Grant of 2015. The authors report no conflicts of interest related to this study.

References

1. Zhou RH. *Resource science of Chinese medicinal materials*. Beijing: China Medical & Pharmaceutical Sciences Press. 1993:202-211.
2. Jang JY, Lee JH, Shin HK, Choi YH, Lee JD, Choi BT. Partially purified Asiasari radix inhibits melanogenesis through extracellular signal-regulated kinase signaling in B16F10 cells. *Int J Mol Med*. 2010;25:287-292.
3. Jeong SH, Lee JE, Jin SH, Ko Y, Park JB. Effects of Asiasari radix on the morphology and viability of mesenchymal stem cells derived from the gingiva. *Mol Med Rep*. 2014;10(6): 3315-3319.
4. Kim HM, Moon YS. Asiasari radix inhibits immunoglobulin E production on experimental models in vitro and in vivo. *Immunopharmacol Immunotoxicol*. 1999;21:469-481.
5. Rho SS, Park SJ, Hwang SL, Lee MH, Kim CD, Lee IH, et al. The hair growth promoting effect of Asiasari radix extract and its molecular regulation. *J Dermatol Sci*. 2005;38:89-97.
6. Jeon HC, Rho EJ, Kim HR, Yun YG. A Study on application of radix Asari main blended prescription from Dongeubogam. *Korean J Oreint med Prescr*. 2004;12(2):57-76.
7. Takara K, Horibe S, Obata Y, Yoshikawa E, Ohnishi N, Yokoyama T. Effects of 19 herbal extracts on the sensitivity to paclitaxel or 5-fluorouracil in HeLa cells. *Biol Pharm Bull*. 2005;28:138-142.
8. Oh SM, Kim J, Lee J, Yi JM, Oh DS, Bang OS, et al. Anticancer potential of an ethanol extract of Asiasari radix against HCT-116 human colon cancer cells in vitro. *Oncol Lett*. 2013;5:305-310.
9. Jeong SH, Kim BB, Lee JE, Ko Y, Park JB. Evaluation of the effects of Angelica dahurica radix on the morphology and viability of mesenchymal stem cells. *Mol Med Rep*. 2015; 12:1556-1560.
10. Jeong SH, Lee JE, Kim BB, Ko Y, Park JB. Evaluation of the effects of Cimicifugae Rhizoma on the morphology and viability of

- mesenchymal stem cells. *Exp Ther Med.* 2015; 10:629-634.
11. Park JB. Restoration of the severely decayed tooth using crown lengthening with simultaneous tooth-preparation. *European journal of dentistry.* 2010;4:197-201.
12. Wang Y, Wang WL, Xie WL, Li LZ, Sun J, Sun WJ, et al. Puerarin stimulates proliferation and differentiation and protects against cell death in human osteoblastic MG-63 cells via ER-dependent MEK/ERK and PI3K/Akt activation. *Phytomedicine.* 2013;20: 787-796.
13. Fournier BP, Ferre FC, Couty L, Lataillade JJ, Gourven M, Naveau A, et al. Multipotent progenitor cells in gingival connective tissue. *Tissue Eng Part A.* 2010;16:2891-2899.
14. Yang H, Gao LN, An Y, Hu CH, Jin F, Zhou J, et al. Comparison of mesenchymal stem cells derived from gingival tissue and periodontal ligament in different incubation conditions. *Biomaterials.* 2013;34: 7033-7047.
15. Su WR, Zhang QZ, Shi SH, Nguyen AL, Le AD. Human gingiva-derived mesenchymal stromal cells attenuate contact hypersensitivity via prostaglandin E2-dependent mechanisms. *Stem Cells.* 2011;29:1849-1860.
16. Zhang Q, Nguyen AL, Shi S, Hill C, Wilder-Smith P, Krasieva TB, et al. Three-dimensional spheroid culture of human gingiva -derived mesenchymal stem cells enhances mitigation of chemotherapy-induced oral mucositis. *Stem Cells Dev.* 2012;21:937-947.
17. Fournier BP, Larjava H, Hakkinen L. Gingiva as a source of stem cells with therapeutic potential. *Stem Cells Dev.* 2013;22:3157-3177.
18. Tomar GB, Srivastava RK, Gupta N, Barhanpurkar AP, Pote ST, Jhaveri HM, et al. Human gingiva-derived mesenchymal stem cells are superior to bone marrow-derived mesenchymal stem cells for cell therapy in regenerative medicine. *Biochem Biophys Res Commun.* 2010; 393:377-383.
19. Chen M, Su W, Lin X, Guo Z, Wang J, Zhang Q, et al. Adoptive transfer of human gingiva-derived mesenchymal stem cells ameliorates collagen-induced arthritis via suppression of Th1 and Th17 cells and enhancement of regulatory T cell differentiation. *Arthritis Rheum.* 2013;65:1181-1193.
20. Zhang Q, Shi S, Liu Y, Uyanne J, Shi Y, Le AD. Mesenchymal stem cells derived from human gingiva are capable of immunomodulatory functions and ameliorate inflammation-related tissue destruction in experimental colitis. *J Immunol.* 2009;183:7787-7798.
21. Zhang QZ, Su WR, Shi SH, Wilder-Smith P, Xiang AP, Wong A, et al. Human gingiva-derived mesenchymal stem cells elicit polarization of m2 macrophages and enhance cutaneous wound healing. *Stem Cells.* 2010; 28:1856-1868.
22. Wang F, Yu M, Yan X, Wen Y, Zeng Q, Yue W, et al. Gingiva-derived mesenchymal stem cell-mediated therapeutic approach for bone tissue regeneration. *Stem Cells Dev.* 2011;20: 2093-2102.
23. Lee SI, Yeo SI, Kim BB, Ko Y, Park JB. Formation of size-controllable spheroids using gingiva-derived stem cells and concave microwells: Morphology and viability tests. *Biomed Rep.* 2016;4:97-101.
24. Santamaria S, Sanchez N, Sanz M, Garcia-Sanz JA. Comparison of periodontal ligament and gingiva-derived mesenchymal stem cells for regenerative therapies. *Clin Oral Investig.* 2016.
25. Strober W. Trypan blue exclusion test of cell viability. *Curr Protoc Immunol.* 2001;Appendix 3:Appendix 3B.
26. Bradford MM. A rapid and sensitive method for the quantitation of microgram quantities of protein utilizing the principle of protein-dye binding. *Anal Biochem.* 1976;72:248-254.

27. Meleti Z, Shapiro IM, Adams CS. Inorganic phosphate induces apoptosis of osteoblast-like cells in culture. *Bone*. 2000;27:359-366.
28. Vande Vannet B, Mohebbian N, Wehrbein H. Toxicity of used orthodontic archwires assessed by three-dimensional cell culture. *Eur J Orthod*. 2006;28:426-432.
29. Sutherland MW, Learmonth BA. The tetrazolium dyes MTS and XTT provide new quantitative assays for superoxide and superoxide dismutase. *Free Radic Res*. 1997;27:283-289.
30. Shi MF, Jiao J, Lu WG, Ye F, Ma D, Dong QG, et al. Identification of cancer stem cell-like cells from human epithelial ovarian carcinoma cell line. *Cell Mol Life Sci*. 2010;67:3915-3925.
31. Zou C, Shen Z. An optimized in vitro assay for screening compounds that stimulate liver cell glucose utilization with low cytotoxicity. *J Pharmacol Toxicol Methods*. 2007;56:58-62.
32. Wagner U, Burkhardt E, Failing K. Evaluation of canine lymphocyte proliferation: comparison of three different colorimetric methods with the 3H-thymidine incorporation assay. *Vet Immunol Immunopathol*. 1999;70:151-159.