

Original Article

Analgesic and Anti-Inflammatory Effect of *Scutellaria Baicalensis*

Joong-Keun Lee, Yun-Kyung Song, Hyung-Ho Lim

Department of Oriental Rehabilitation Medicine, College of Oriental Medicine,
Kyungwon University

Backgrounds : *Scutellaria baicalensis* has been used as a medicinal plant to treat various disease conditions accompanying inflammatory response and oxidative stress.

Objectives : The aim of this study is to evaluate the effects of *Scutellaria baicalensis* against inflammatory, pain and edema

Methods : *In vitro*, the effects of *Scutellaria baicalensis* against lipopolysaccharide-induced inflammation were investigated in mouse BV2 microglial cells. *In vivo*, the effects of *Scutellaria baicalensis* on acetic acid-induced writhing response, carrageenan-induced edema and the plantar test (nociceptive thermal stimulation) were investigated using rats and mice.

Results : The present results showed that pre-treatment with the aqueous extract of *Scutellaria baicalensis* suppressed the lipopolysaccharide-stimulated cyclooxygenase-2 expressions in mouse BV2 microglial cells. The aqueous extract of *Scutellaria baicalensis* inhibited acetic acid-induced abdominal pain in mice and also reduced thermal pain in rats. However, no significant inhibition on carrageenan-induced edema in rats.

Conclusions : The present study showed that *Scutellaria baicalensis* possesses anti-inflammatory and analgesic effects.

Key Words : *Scutellaria baicalensis*, analgesic, anti-inflammatory

Introduction

Scutellaria baicalensis Georgi (Labiatae), called 'Hwang-Gum' in Korea and as 'Huang-Qin' in China, is one of the most widely used herbal medicines against bacterial infection of the respiratory and gastrointestinal tracts and various inflammatory diseases^{1,2)}.

Prostaglandins (PGs) are key inflammatory mediators that are converted from arachidonic

acid by cyclooxygenase (COX). There are two isoforms of COX: cyclooxygenase-1 (COX-1) is constitutively expressed in nearly all tissues and provides PGs to maintain physiological functions such as cytoprotection of the stomach and regulation of renal blood flow^{3,4)}. In contrast, cyclooxygenase-2 (COX-2) is inducible in the immune cells such as macrophages and synoviocytes in response to infection, injury or other stresses and COX-2 produces excessive amount of PGs that sensitize nociceptors and induce inflammatory states^{5,6)}. Prostaglandin E2 (PGE2) is overproduced at sites of inflammation as a result of the activation of inducible COX-2^{7,8)}.

The writhing test is a well-known animal test for the model of inflammatory and visceral pain.

• received : 19 November 2007

• received in revised form : 20 November 2007

• accepted : 5 December 2007

• Correspondence to : Hyung-Ho Lim

Department of Oriental Rehabilitation Medicine, Seoul
Oriental Hospital Kyungwon University, 20-8 Songpa-Dong,
Songpa- Gu, Seoul Korea

(Tel : +82-2-425-3456 / Fax : +82-2-425-3560,

E-mail : omdlimhh@chol.com)

It is useful test for studying anti-nociceptive effect. Carrageenan induced local inflammation is commonly used to evaluate analgesic effect. Therefore, carrageenan induced local inflammation (paw edema) is a useful model to assess the contribution of mediators involved in vascular changes associated with acute inflammation⁹. Plantar test (Hargreaves method) is used to determine thermal pain threshold by exposing the animals to heat¹⁰.

The effect of herbal medicine on inflammatory disease and pain is widely studied. For example, anti-inflammatory of *Corydalis turtschaninovia*¹¹⁻¹⁴, *Clematis mandshurica*^{15,16} are generally evaluated. *Scutellaria baicalensis* is also studied about acute renal failure¹⁷, hepatocytes cytotoxicity¹⁸, ischemia-reperfusion injuries of hearts¹⁹, neuroprotective of the brain ischemia²⁰ and so on. But, anti-inflammatory and analgesic actions of the *Scutellaria baicalensis* have not been clarified yet. In the present study, we investigated the effects of *Scutellaria baicalensis* against LPS-stimulated expressions of COX-1 and COX-2 in mouse BV2 microglial cells by using Western blot analysis in mouse BV2 microglial cells, *in vitro*. Moreover, the anti-nociceptive and anti-inflammatory effects of *Scutellaria baicalensis* using several experimental pain models, acetic acid-induced writhing response, carrageenan-induced for edema and response on thermal pain were also evaluated, *in vivo*.

Materials and Methods

1. Preparation of the aqueous extract of *Scutellaria baicalensis*

To obtain the aqueous extract of *Scutellaria baicalensis*, 100 g of *Scutellaria baicalensis* was

added to distilled water and extraction was performed by heating at 90°C for 2 h, concentrating with a rotary evaporator and lyophilization (Eyela, Tokyo, Japan). The resulting powder, weighing 20.58 g, was dissolved in saline solution and filtered through a 0.22 μm syringe before use.

2. *In vitro* experiments

1) Reagents

The *Scutellaria baicalensis* was purchased from Oriental Medicine Hospital, Kyungwon University (Seoul, Korea). Rats and mice were purchased from Oriental Bio Co (Kyunggido, Korea).

2) Cell culture

Mouse BV2 macroglia cells were cultured in Dulbecco's modified Eagle's Medium (DMEM; Gibco BRL, Grand Island, NY, USA) supplemented with 10 % heat-inactivated fetal bovine serum (FBS; Gibco BRL, Grand Island, NY, USA) at 37°C in 5 % CO₂-95 % O₂ in a humidified cell incubator. The cells were plated onto culture dishes at a density of 2×10^4 cells/cm² in culture dish, 24 h prior to drug treatments.

3) MTT cytotoxicity assay

Mouse BV2 macroglia cells were grown in a final volume of 100 μl culture medium per well in 96-well plates. In order to determine the cytotoxicity of *Scutellaria baicalensis*, the cells were treated with *Scutellaria baicalensis* at concentrations of 1 $\mu\text{g/ml}$, 10 $\mu\text{g/ml}$, 100 $\mu\text{g/ml}$, 1,000 $\mu\text{g/ml}$ and 10,000 $\mu\text{g/ml}$ for 24 h. Cultures of the cells of the control group were left untreated. After adding 10 μl of the MTT labeling reagent containing 5 mg/ml 3-(4,5-dimethyl thiazol-2-yl)-2,5-diphenyl tetrazolium bromide in phos-

phate-buffered saline to each well, the plates were incubated for 2 h. Solubilization solution 100 μl containing 10 % sodium dodecyl sulfate in 0.01 M hydrochloric acid was added to each well and the cells were incubated for another 12 h. The absorbance was then measured with a microtiter plate reader (Bio-Tek, Winooski, VT, USA) at a test wavelength of 595 nm with a reference wavelength of 690 nm. The optical density (O.D.) was calculated as the difference between the absorbance at the reference wavelength and that observed at the test wavelength. Percent viability was calculated as (O.D. of drug-treated sample/control O.D.) \times 100.

4) Western blot analysis

Mouse BV2 microglial cells were lysed in a ice-cold whole cell lysate buffer containing 50 mM HEPES (pH 7.5), 150 mM NaCl, 10 % glycerol, 1 % Triton X-100, 1.5 mM magnesium chloride hexahydrate, 1 mM ethyleneglycol-bis-(β -aminoethyl ether)-N,N'-tetraacetic acid (EGTA), 1 mM phenylmethylsulfonyl fluoride (PMSF), 2 $\mu\text{g}/\text{ml}$ leupeptin, 1 $\mu\text{g}/\text{ml}$ pepstatin, 1 mM sodium ortho vanadate and 100 mM sodium fluoride and the mixture were incubated 30 min at 4°C. Cell debris was removed by microcentrifugation, followed by quick freezing of the supernatant. The protein concentration was measured using a Bio-Rad colorimetric protein assay kit (Bio-Rad, Hercules, CA, USA). Protein of 40 μg was separated on SDS-polyacrylamide gels and transferred onto a nitrocellulose membrane (Schleicher & Schuell GmbH, Dassel, Germany). Goat COX-1 antibody (1:1000; Santa Cruz Biotech, CA, USA) and Goat COX-2 antibody (1:1000; Santa Cruz Biotech, CA, USA) were used as a primary antibody. Horseradish peroxidase-conjugated anti-goat antibody (1:4000; Santa Cruz Biotech

for COX-1 and COX-2 was used as a secondary antibody. Band detection was performed using the enhanced chemiluminescence (ECL) detection system (Amersham Pharmacia Biothech GmbH, Freiburg, Germany).

3. *In vivo* experiments

1) Animals and treatments

Male ICR mice weighing 28-30 g and male Sprague-Dawley rats weighing 150-160 g were used for the experiments. The experimental procedures were performed in accordance with the animal care guidelines of the National Institute of Health (NIH) and the Korean Academy of Medical Sciences. The animals were housed under laboratory conditions at a controlled temperature (20 \pm 2°C) and maintained under light-dark cycles, each consisting of 12 h of light and 12 h of darkness (lighting from 07:00 to 19:00 h) with food and water made available *adlibitum*.

2) Acetic acid-induced writhing response in mice

Male ICR mice were divided into five groups: the control group, the acetic acid injection group, the acetic acid injection and 50 mg/kg *Scutellaria baicalensis*-treated group, the acetic acid injection and 100 mg/kg *Scutellaria baicalensis*-treated group and the acetic acid injection and 200 mg/kg *Scutellaria baicalensis*-treated group (n = 10 in each group). *Scutellaria baicalensis* and control vehicle were orally administered 1 h before the acetic acid injection. Then, the mice received intraperitoneally with 0.15 ml of 1.0 % acetic acid as an irritant stimulus and placed in an individual plastic cage (20 \times 30 \times 12 cm high) for observation. The number of writhes was counted for 30 min after acetic acid injection.

3) Carrageenan-induced edema in rats

The volume of the paw edema in male Sprague-Dawley rats was measured in each animal using a plethysmometer (Ugo Basile, Italy) with a precision of two decimal places. To induce edema in the experimental animals, a single subplantar injection of carrageenan (1 %, 0.05 ml Sigma Chemical Co., St. Louis, MO, USA) was given to each animal and the animals of the control group received injections of equivalent doses of normal saline²¹⁾ as a same method.

Animals of *Scutellaria baicalensis*-treated groups received orally with 1 ml of the aqueous extract of *Scutellaria baicalensis* at the respective doses before 1 h of carrageenan injection and those of the control group and the carrageenan-induced edema group received equivalent amount of drinking water before 1 h of carrageenan injection. The paw volume was measured immediately before, 1 h, 2 h, 3 h, 4 h and 5 h after carrageenan injection. The animals were divided into five groups: the control group, the carrageenan-induced edema group, the carrageenan-induced edema and 100 mg/kg *Scutellaria baicalensis*-treated group, the carrageenan-induced edema and 200 mg/kg *Scutellaria baicalensis*-treated group and the carrageenan-induced edema and 400 mg/kg *Scutellaria baicalensis*-treated group (n = 10 in each group). The percentage of edema was calculated as follows:

$$\text{Percentage of edema (\%)} = (V_t - V_n) / V_n \times 100$$

V_t = The paw volume of each time after the injection of carrageenan

V_n = The paw volume before the injection of carrageenan

4) Plantar test (Hargreaves's method)

To assess nociceptive responses to thermal stimuli, paw withdrawal latency of male Sprague-

Dawley rats was tested using the procedure previously described by Hargreaves et al¹⁰⁾. The center of a focused beam of radiant heat was applied to the plantar surface of the hind paw in rats and the withdrawal latency time was recorded. The infra-red intensity of the heat stimulus was 60 and stimulation was adjusted so that the baseline latency was 6 sec and 20 sec cut-off time was imposed to avoid tissue damage. Three minutes were allowed for the next test. The animals in the *Scutellaria baicalensis*-treated groups received orally with 1 ml of the aqueous extract of *Scutellaria baicalensis* at the respective doses before 1 h of test and those of the control group received equivalent amount of saline before 1 h of test. The withdrawal latency time was measured immediately before, 1 h and 2 h after drug administration. The animals were divided into four groups: thermal stimulation-induced nociception and drinking water-treated group, thermal stimulation-induced nociception and 50 mg/kg *Scutellaria baicalensis*-treated group, thermal stimulation-induced nociception and 100 mg/kg *Scutellaria baicalensis*-treated group and thermal stimulation-induced nociception and 200 mg/kg *Scutellaria baicalensis*-treated group (n = 6 in each group).

4. Data analysis

Results are presented as the mean \pm standard error means (SEM). Statistical analysis was performed using one-way ANOVA followed by Duncan *post-hoc* test. Duncan's Test can be used to determine the significant differences between group means in an analysis of variance setting. Duncan's test is based on the range statistic for a detailed discussion of different post hoc tests. Difference was considered significant at $p < 0.05$.

Results

1. Cell viability of *Scutellaria baicalensis* on BV2 microglia cells

In order to assess the cytotoxic effect of the aqueous extract of *Scutellaria baicalensis* on BV2 microglia cells, the cells were cultured with the aqueous extract of *Scutellaria baicalensis* at final concentrations of 0.001, 0.01, 0.1, 1 and 10 mg/ml for 24 h and MTT assays were then carried out. The cells cultured in *Scutellaria baicalensis*-free media were used as the control. The viability of cells incubated with *Scutellaria baicalensis* at concentrations of 0.001, 0.01, 0.1, 1 and 10 mg/ml for 24 h was 105.10 ± 0.94 , 103.83 ± 1.70 , 106.51 ± 1.71 , 77.31 ± 2.06 and 30.07 ± 0.47 % of the control value, respectively.

The present results show that the *Scutellaria baicalensis* exerted no significant cytotoxicity until it was at a concentration of 0.1 mg/ml. However, a high concentration (1 mg/ml and 10 mg/ml) of *Scutellaria baicalensis* reduced cell viability. Then, we used the aqueous extract of *Scutellaria baicalensis* at concentration of 0.01 and 0.1 mg/ml for the next experiment (Fig. 1).

2. Effect of *Scutellaria baicalensis* on protein expression of COX-1 and COX-2

The level of COX-1 protein was 1.06 ± 0.02 following a treatment with 1 $\mu\text{g/ml}$ LPS for 24 h. The level of COX-1 protein was 0.97 ± 0.01 , 0.86 ± 0.04 and 0.80 ± 0.05 in the cells pre-treated with the aqueous extract of *Scutellaria baicalensis* at 0.01 mg/ml, 0.1 mg/ml and 500 μM acetylsalicylic acid (ASA), one hour before LPS treatment.

The level of COX-2 protein was markedly increased to 5.28 ± 0.81 following a treatment with 1 $\mu\text{g/ml}$ LPS for 24 h. The level of COX-2 protein was decreased to 3.64 ± 0.58 , 1.27 ± 0.16 and 1.49 ± 0.08 in the cells pre-treated the aqueous extract of *Scutellaria baicalensis* at 0.01 mg/ml, 0.1 mg/ml and 500 μM ASA one hour before LPS treatment.

The present results show that LPS enhanced COX-2 protein expression in mouse BV2 microglia cells and the aqueous extract of *Scutellaria baicalensis* suppress LPS-induced COX-2 protein expression. However, LPS and the aqueous extract of *Scutellaria baicalensis* exerted no significant effect on the expression of




Fig. 1. Cell viability of *Scutellaria baicalensis* on BV2 microglia cells. (A) Control group, (B) 0.001 mg/ml *Scutellaria baicalensis*-treated group, (C) 0.01 mg/ml *Scutellaria baicalensis*-treated group, (D) 0.1 mg/ml *Scutellaria baicalensis*-treated group, (E) 1 mg/ml *Scutellaria baicalensis*-treated group, (F) 10 mg/ml *Scutellaria baicalensis*-treated group. * represents $p < 0.05$ compared to the control group.




Fig. 2. Results of Western blot analysis of the protein levels of COX-1 and COX-2. (A) Control, (B) LPS-treated group, (C) LPS- and 0.01 mg/ml *Scutellaria baicalensis*-treated group, (D) LPS- and 0.1 mg/ml *Scutellaria baicalensis*-treated group, (E) LPS- and 500 μM ASA-treated group. Actin was used as the internal control. * represents $p < 0.05$ compared to the control group. # represents $p < 0.05$ compared to the LPS-treated group.

COX-1 protein, except the aqueous extract of *Scutellaria baicalensis* at concentration of 0.1 mg/ml (Fig. 2).

3. Effect of *Scutellaria baicalensis* on acetic acid-induced writhing response in mice

The number of the writhing response in the control group was 0.25 ± 0.25 . The number of writhing response in the acetic acid injection group was 39.88 ± 8.18 . The number of writhing response in the acetic acid injection and *Scutellaria baicalensis* (50, 100 and 200 mg/kg)-treated group was 27.33 ± 1.61 , 29.33 ± 6.10




Fig. 3. Effect of *Scutellaria baicalensis* on the number of writhing response. (A) control group, (B) 1% acetic acid-induced writhing response group, (C) 1% acetic acid-induced writhing response and 50 mg/kg *Scutellaria baicalensis*-treated group, (D) 1% acetic acid-induced writhing response and 100 mg/kg *Scutellaria baicalensis*-treated group, (E) 1% acetic acid-induced writhing response and 200 mg/kg *Scutellaria baicalensis*-treated group. * represents $p < 0.05$ compared to the control group. # represents $p < 0.05$ compared to the 1% acetic acid-induced writhing response group.

and 25.63 ± 3.04 .

The present results showed that acetic acid injection into the abdominal cavity induced writhing response. The acetic acid injection and *Scutellaria baicalensis*-treated group suppressed acetic acid-induced writhing response (Fig. 3).

4. Effect of *Scutellaria baicalensis* on the volume of carrageenan-induced paw edema

After 1 h, paw volume in the control group was 0.00 ± 0.00 %. Paw volume in the 1 % carrageenan-induced edema group was increased to 5.96 ± 0.28 %. Paw volume of 1 % carrageenan-induced edema and *Scutellaria baicalensis*-treated groups at concentrations of 100, 200 and 400 mg/kg was 4.68 ± 1.79 , 9.40 ± 1.52 and 7.87 ± 2.17 %.

After 2 h, paw volume in the control group was 0.00 ± 0.00 %. Paw volume in the 1 % carrageenan-induced edema group was increased to 17.57 ± 2.11 %. Paw volume in the 1 %

carrageenan-induced edema and *Scutellaria baicalensis*-treated groups at concentrations of 100, 200 and 400 mg/kg was 14.61 ± 1.23 , 22.51 ± 3.97 and 14.92 ± 2.26 %.

After 3 h, paw volume in the control group was 0.00 ± 0.00 %. Paw volume in the 1 % carrageenan-induced edema group was increased to 24.21 ± 2.21 %. Paw volume in the 1 % carrageenan-induced edema and *Scutellaria baicalensis*-treated groups at concentrations of 100, 200 and 400 mg/kg was 27.86 ± 1.77 , 28.00 ± 4.78 and 20.67 ± 2.52 %.

After 4 h, paw volume in the control group was 0.00 ± 0.00 %. Paw volume in the 1 % carrageenan-induced edema group was increased to 26.20 ± 3.39 %. Paw volume in the 1 % carrageenan-induced edema and *Scutellaria baicalensis*-treated groups at concentrations of 100, 200 and 400 mg/kg was 21.80 ± 3.21 , 30.38 ± 3.74 and 20.67 ± 2.52 %.

After 5 h, paw volume in the control group was 0.00 ± 0.00 %. Paw volume in the 1 %




Fig. 4. Effect of *Scutellaria baicalensis* in response to doses and time on carrageenan-induced paw edema. (□) control group, (■) 1 % carrageenan-induced edema group, (○) 1 % carrageenan-induced edema and 100 mg/kg *Scutellaria baicalensis*-treated group, (●) 1 % carrageenan-induced edema and 200 mg/kg *Scutellaria baicalensis*-treated group, (△) 1 % carrageenan-induced edema and 400 mg/kg *Scutellaria baicalensis*-treated group.

carrageenan-induced edema group was increased to $15.71 \pm 4.56\%$. Paw volume in the 1% carrageenan-induced edema and *Scutellaria baicalensis*-treated groups at concentrations of 100, 200 and 400 mg/kg was 8.24 ± 2.95 , 14.83 ± 5.51 and $21.19 \pm 6.63\%$.

The present results showed that the paw volume in the control group was maintained at constant level and that paw volume in the carrageenan-induced edema group was increased as time-dependently during 4 h.

First of all, the concentrations of *Scutellaria baicalensis*-treated groups are 50, 100, 200 mg/kg. However, carrageenan-induced edema and *Scutellaria baicalensis*-treated groups exerted no significant inhibition on carrageenan-induced paw edema. And this time, the concentrations of *Scutellaria baicalensis*-treated groups changed 100, 200, 400 mg/kg, but the result was also no significant inhibition (Fig. 4).

5. Effect of *Scutellaria baicalensis* on the plantar test (nociceptive thermal stimulation)

After 1 h, paw withdrawal threshold in the pre-treated value was considered as 1.00. The withdrawal latency in the thermal stimulation-induced nociception and drinking water-treated group was 1.17 ± 0.17 . The withdrawal latency in the thermal stimulation-induced nociception and 50 mg/kg *Scutellaria baicalensis*-treated group was 1.39 ± 0.13 . The withdrawal latency in the thermal stimulation-induced nociception and 100 mg/kg *Scutellaria baicalensis*-treated group was 1.13 ± 0.16 . The withdrawal latency in the thermal stimulation-induced nociception and 200 mg/kg *Scutellaria baicalensis*-treated group was 1.28 ± 0.15 . After 1 h, the withdrawal latency of thermal stimulation-induced nociception was not significantly different in the animals treated with *Scutellaria baicalensis*.

After 2 h, the withdrawal latency in the thermal stimulation-induced nociception and drinking water-treated group was 0.96 ± 0.09 . The withdrawal latency in the thermal stimulation-induced nociception and 50 mg/kg *Scutellaria baicalensis*-treated group was 1.22 ± 0.15 . The




Fig. 5. Effect of *Scutellaria baicalensis* in response to doses and time on thermal pain. (□) thermal stimulation-induced nociception and drinking water-treated group (control), (■) thermal stimulation-induced nociception and 50 mg/kg *Scutellaria baicalensis*-treated group, (○) thermal stimulation-induced nociception and 100 mg/kg *Scutellaria baicalensis*-treated group, (●) thermal stimulation-induced nociception and 200 mg/kg *Scutellaria baicalensis*-treated group. * represents $p < 0.05$ compared to the control group.

withdrawal latency in the thermal stimulation-induced nociception and 100 mg/kg *Scutellaria baicalensis*-treated group was 1.03 ± 0.08 . The withdrawal latency in the thermal stimulation-induced nociception and 200 mg/kg *Scutellaria baicalensis*-treated group was 1.36 ± 0.16 . The present results showed that the withdrawal latency in the thermal stimulation-induced nociception and 200 mg/kg *Scutellaria baicalensis*-treated group was increased after 120 min. *Scutellaria baicalensis* showed significant inhibitory effect on thermal stimulation-induced nociception (Fig. 5).

Discussion

Scutellaria baicalensis is known to have the anti-bacterial activity against staphylococci, cholera, typhoid, paratyphoid, dysentery, diphtheria, hemolytic streptococci, Escherichia coli, pneumococci and spirochaeta^{1,2)} and it also exerts anti-viral effect against influenza virus²⁾. The properties of *Scutellaria baicalensis* is 'bitter and cold'. It is credited with 'damp-heat'-clearing, 'heat'-lower, detoxicant and anti-inflammatory actions and is known to prevent abnormal fetal movements¹⁾. It has been used for the treatment of fever, cough, hemoptysis, jaundice, hepatitis, dysentery, acute conjunctivitis, carbuncle and furuncle¹⁾.

Scutellaria baicalensis is known to contain a number of flavone derivatives²²⁾. The first flavone isolated from its root was wogonin. Wogonin is present only in small amounts in the root; the flavone glycoside named baicalin. Acid hydrolysis of baicalin yields glucuronic acid and a flavone aglycone named baicalein. Anti-inflammatory effects of these three major constituents-baicalin, baicalein and wogonin-are

well documented²³⁻²⁵⁾. Baicalin, baicalein and wogonin were found to inhibit acetic acid-induced increase in vascular permeability in mice and to reduce acute paw edema in rats induced by compound 48/80²³⁾. They also suppressed development of secondary lesion in adjuvant-induced arthritis in rats²³⁾. Baicalein and wogonin have been shown to inhibit lipopolysaccharide (LPS)-induced nitric oxide (NO) production in macrophages²⁵⁾. Flavonoids can inhibit 5-lipoxygenase and cyclooxygenase activities²⁶⁾ and act as phospholipase A2 inhibitors²⁷⁾. These evidences indicate that flavonols can modulate the prostanoid biosynthetic pathway. Arachidonic acid which is accumulated in the membrane lipid, can be selectively released from the phospholipids pool by chemical or mechanical stimulation and is subsequently converted to prostaglandins (PGs) by two enzymes, COX-1 and COX-2²⁸⁾. COX-2 is primarily responsible for PGs produced in inflammation and COX-1 for PGs involved in normal homeostasis. In this regard, COX-2 is up-regulated in the air pouch and catalyzes the production of large amounts of PGE2²⁹⁾. The up-regulation of COX-2 associated with the increase of PGE2 may be major event in acetic acid-induced writhing response and carrageenan-induced inflammation. Alternatively, the decrease of PGE2 may be induced by the inhibition of the release of TNF- α , because stimulation of macrophages/monocytes, fibroblasts and epithelial cells with cytokines such as IL-1 and TNF- α leads to PGE2 production²⁸⁾. In the present results, the aqueous extract of *Scutellaria baicalensis* suppresses LPS-induced COX-2 protein expression in mouse BV2 microglia cells.

The analgesic effect of *Scutellaria baicalensis*

was evaluated by acetic acid-induced writhing response. In acetic acid-induced abdominal writhing which is the visceral pain model, the processor releases arachidonic acid via cyclooxygenase and prostaglandin biosynthesis plays a role in the nociceptive mechanism³⁰. In the present results, pre-treatment with the aqueous extract of *Scutellaria baicalensis* revealed an analgesic effect on acetic acid-induced writhing response in mice.

The carrageenan test was selected because of its sensitivity in detecting orally active anti-inflammatory agents particularly in the acute phase of inflammatory⁹. The intraplantar injection of carrageenan in rats leads to paw edema. In the present results, however, *Scutellaria baicalensis* exerted no significant inhibition on carrageenan-induced paw edema.

Thermal hyperalgesia can be explained by central convergence of afferents from deep tissues and the skin³¹. In the present results, administration of *Scutellaria baicalensis* suppressed the thermal stimulation-induced nociception.

Here in this study, we have demonstrated that the aqueous extract of *Scutellaria baicalensis* have analgesic and anti-inflammatory activities and could be the reason for its wide use in traditional medicine to treat different types of pain. Although pharmacological activities have been carried out, the detailed mechanisms of actions on each respective effect should be further investigated.

References

1. Chang HM, But PP. Pharmacology and Applications of Chineses Materia Medica, vol. 1. Singapore:World Scientific. 1986:1022.
2. Huang KC. The Pharmacology of Chinese Herbs. 1st ed. Boca Raton:CRC Press. 1993: 291.
3. Vane JR, Botting RM. Mechanism of action of nonsteroidal anti-inflammatory drugs. Am J Med. 1998; 104:2-S8.
4. Vane JR, Bakhle YS, Botting RM. Cyclooxygenases 1 and 2. Annu Rev Pharmacol. Toxicol. 1998; 38:97-120.
5. Duval DL, Miller DR, Collier J, Billings RE. Characterization of hepatic NO synthase: identification as the cytokine-inducible form primarily regulated by oxidants. Mol Pharmacol. 1996; 50:277-84.
6. Marletta MA. Nitric oxide synthase structure and mechanism. J Biol Chem. 1993; 268: 12231-4.
7. Fu JY, Masferrer JL, Seibert K, Raz A, Needleman P. The induction and suppression of prostaglandin H2 synthase (cyclooxygenase) in human monocytes. J Biol Chem. 1990; 265:16737-40.
8. Picot D, Loll PJ, Garavito RM. The X-ray crystal structure of the membrane protein prostaglandin H2 synthase-1. Nature. 1994; 367:243-9.
9. Di Rosa M, Giroud JP, Willoughby DA. Studies on the mediators of the acute inflammatory response induced in rats in different sites by carrageenan and turpentine. J Pathol. 1971; 104(1):15-29.
10. Hargreaves K, Dubner R, Brown F, Flores C, Joris J. A new and sensitive method for measuring thermal nociception in cutaneous hyperalgesia. Pain. 1988; 32(1):77-88.
11. Kim IK, Lee TH, Kim CJ, Lee CY. Effects of Herbal Bath on Acetic Acid-induced Somato-visceral Pain in Mice. The Journal of Oriental Physiology & Pathology. 2006; 20(3):642-50.
12. Kim BS, Kim YS. Effect of *Corydalis Tuber* · *Paeoniae Radix Rubra* · *Carthami Flos* Complex Herbal Acupuncture on Inflammation.

- mation and Liver in Rats with Arthritis induced by Freund's Complete Adjuvant. The Journal of Herbology. 2004; 19(1):95-102.
13. Ahn HS, Park YB, Kang SK. Effects of the *Corydalis Tuber* Aqua-acupuncture on the Pain and Convulsion in Mice and Gastric Ulcer in Rats. The Journal of Korean Acupuncture & Moxibustion Society. 1994; 11(1) :99-111.
 14. Ahn CJ, Ahn BC, Park DS. The analgesic effect of aqua-acupuncture with *Corydalis Tuber*, *Olibanum Mastrix* & *Myrrha*. The Journal of Korean Acupuncture & Moxibustion Society. 1994; 11(1):145-58.
 15. Cho SW, Kim KS. Mechanism of Herbal Acupuncture of *Clematis Mandshurica Maxim*. Water Extract by Stimulation of Sinsu (BL 23) Loci Subcutaneously as Dual Inhibitor of Proinflammatory Cytokines on Adjuvant Arthritis in Rats. Korean Journal of Oriental Medicine. 2006; 27(4):182-90.
 16. Yang KJ, Kim SJ, Seo IB, Park SK, Kim JS, Seo JC, Choi SM, Lee HJ, Kim EH. Effects of *Clematis mandshurica* Rupr. on Nitric Oxide Synthase in the Periaqueductal Gray of Collagenase-induced Rat Osteoarthritis Model. The Korean Journal of Meridian & Acupoint. 2006; 22(4):109-16.
 17. Kim SC, Lee SD, Song SH, Koh PO, Kang CB, Kim JS, Kim CH. Effect of *Scutellaria Baicalensis* Georgi. Extract on Cisplatin-Induced Acute Renal Failure in Rabbits. The Journal of Veterinary Clinics. 2007; 24(3): 392-9.
 18. Byun BH, Seo BI. Molecular Cell Biological Study on Rat Hepatocytes Cytotoxicity and DNA Synthesis Effects of *Scutellariae Baicalensis* Radix Herbal acupuncture Solution (SBRHA). The Journal of Herbology. 2004; 19(1):35-40.
 19. Kim CH, Jia H, Kim SH, Moon HJ, Lee JS. The Protective Effect of *Scutellaria Baicalensis* Georgi on Ischemia-reperfusion Injuries of Rat Hearts. The Korean Journal of Laboratory Animal Science. 2004; 20(4): 357-62.
 20. Lee BC, Leem KH, Kim YO, Kim SY, Ahn DK, Park HK, Kim HC. Neuroprotective Effects of *Scutellaria Radix* on the Brain Ischemia induced by Four -Vessel Occlusion in Rats. The Journal of Herbology. 1999; 14(2):89-96
 21. Sluka KA, Chandran P. Enhanced reduction in hyperalgesia by combined administration of clonidine and TENS. Pain. 2002; 100 (1-2):183-90.
 22. Zhang YY, Don HY, Guo YZ, Ageta H, Harigaya Y, Onda M, Hashimoto K, Ikeya Y, Okada M, Maruno M. Comparative study of *Scutellaria planipes* and *Scutellaria baicalensis*. Biomedical Chromatography. 1998 12:31-3.
 23. Kubo M, Matsuda H, Tanaka M, Kimura Y, Okuda H, Higashino M, Tani T, Namba K, Arichi S. Studies on *Scutellariae radix*. VII. Anti-arthritic and anti-inflammatory actions of methanolic extract and flavonoid components from *Scutellariae radix*. Chemical and Pharmaceutical Bulletin (Tokyo). 1984; 32: 2724-9.
 24. Sekiya K, Okuda H. Selective inhibition of platelet lipooxygenase by baicalein. Biochemical and Biophysical Research Communications. 1982; 105:1090-5.
 25. Wakabayashi I. Inhibitory effects of baicalein and wogonin on lipopolysaccharide-induced nitric oxide production in macrophages. Pharmacology and Toxicology. 1999; 84: 288-91.
 26. Loughton MJ, Evans PJ, Moroney MA, Houlton JR, Halliwell B. Inhibition of mammalian 5-lipoxygenase and cyclo-oxygenase by flavonoids and phenolic dietary additives.

- Relationship to antioxidant activity and to iron ion-reducing ability. *Biochemical Pharmacology*. 1999; 42:1673-81.
27. Lindahl M, Tagesson C. Flavonoids as phospholipase A2 inhibitors: importance of their structure for selective inhibition of group II phospholipase A2. *Inflammation*. 1997; 21:347-56.
 28. Griffiths RJ. Prostaglandins and inflammation. In: Gallin JI, Snyderman R. *Inflammation. Basic Principle and Clinical Correlates*. 3rd ed. Philadelphia:Lippincott Williams and Wilkins Press. 1999:349-60.
 29. Masferrer JL, Zweifel BS, Manning PT, Hauser SD, Leahy KM, Smith WG, Isakson PC, Seibert K. Selective inhibition of inducible cyclooxygenase 2 *in vivo* is anti-inflammatory and nonulcerogenic. *Proceedings of National Academy of Science of the United States of America*. 1994; 91:3228-32.
 30. Franzotti EM, Santos CV, Rodrigues HM, Mourao RHandrade MR, Antonioli AR. Anti-inflammatory, analgesic activity and acute toxicity of *Sida cordifolia* L. (Malvaceae). *J Ethnopharmacol*. 2000; 72(1-2): 273-7.
 31. Light A, Perl E. Unmyelinated afferent fibers are not only for pain anymore. 1st ed. *J Comp Neurol*. 2003; 461:137-9.