Original Article

Development of Standardized Predictive Models for Traditional Korean Medical Diagnostic Pattern Identification in Stroke Subjects: A Hospital-based Multi-center Trial

Woo-Sang Jung, Seung-Yeon Cho, Seong-Uk Park, Sang-Kwan Moon,
Jung-Mi Park, Chang-Nam Ko, Ki-Ho Cho, Seungwon Kwon

Department of Cardiology and Neurology, College of Korean Medicine, Kyung Hee University

Objectives: To develop a standardized diagnostic pattern identification equation for stroke patients, our group conducted a study to derive the predictive logistic equations. However, the sample size was relatively small. In the current study, we aimed to derive new predictive logistic equations for each diagnostic pattern using an expanded number of subjects.

Methods: This study was a hospital-based multi-center trial recruited stroke patients within 30 days of symptom onset. Patients' general information, and the variables related to diagnostic pattern identification were measured. The diagnostic pattern of each patient was identified independently by two Korean Medicine Doctors. To derive a predictive model for pattern identification, binary logistic regression analysis was applied.

Results: Among the 1,251 patients, 385 patients (30.8%) had the Fire Heat Pattern, 460 patients (36.8%) the Phlegm Dampness Pattern, 212 patients (16.9%) the Qi Deficiency Pattern, and 194 patients (15.5%) the Yin Deficiency Pattern. After the regression analysis, the predictive logistic equations for each pattern were determined.

Conclusion: The predictive equations for Fire Heat, Phlegm Dampness, Qi Deficiency, and Yin Deficiency would be useful to determine individual stroke patients' pattern identification in the clinical setting. However, further studies using objective measurements are necessary to validate these data.

Key Words : Fire Heat; Phlegm Dampness; Qi Deficiency; Yin Deficiency; Pattern identification; Stroke

Introduction

Stroke can be fatal or lead to irreversible neurological deficits such as hemiplegia, problems with speech or comprehension, or sense dysfunction¹⁾. To prevent occurrence of irreversible neurological deficits and secondary stroke, several acute stroke management therapies are employed, and include

treatment with recombinant tissue plasminogen activator (rtPA), a protein involved in the breakdown of blood clots^{2,3)}, antiplatelet agents⁴⁾ in ischemic stroke and antihypertensive therapy^{5,6)}, aspiration surgery⁶⁾ in hemorrhagic stroke. However, these therapies have drawbacks, such as the short time window for administration of rtPA and the adverse effects, including bleeding,

• Received : 26 October 2019

• Revised : 19 November 2019

• Accepted : 22 November 2019

• Correspondence to : Seungwon Kwon

Department of Cardiology and Neurology, College of Korean Medicine, Kyung Hee University

23 KyunHee-daero, Dongdaemun-gu, Seoul 02447, Korea

Tel : +82-2-958-9127, Fax : +82-2-958-9132, E-mail : kkokkottung@hanmail.net

)

induced by rtPA or antiplatelets²⁻⁴⁾.

The alternative and complementary medicine therapies have received recent attention to make up for those limitations of western medicine⁷⁾. Previous studies have suggested that therapies such as herbal medicine, acupuncture, electroacupuncture, moxibustion and pharmacopuncture are effective in various fields of stroke therapy such as acute of small vessel occlusion (SVO)^{11,12)}, post-stroke spasticity¹³⁾, motor recovery after stroke¹⁴⁻¹⁶⁾, post-stroke aphasia¹⁷⁾ and shoulder pain induced by hemiplegia¹⁸⁾.

Pattern-identification is a unique diagnostic system in Korean Medicine (KM), Traditional Chinese Medicine and Japanese Kampo medicine. It is an approach to investigate the causative factors, location, and prognosis of diseases according to correlation of all four examinations such as inspection, inquiry, listening and smelling examination, and palpation¹⁹⁾. According to these pattern-identifications, future therapeutic plans are set. Therefore, proper diagnostic pattern-identification is essential to promote the beneficial effects of therapies. To enhance the objectivity of diagnostic pattern-identification, several studies have been conducted in the East Asian region²⁰⁻²⁷⁾. However, most of these studies did not suggest objective coefficient values that could indicate the importance of variables for determining a diagnostic pattern. Furthermore, most²¹⁻²⁶⁾ were based on a very small sample size. Most importantly, the subjects of most of these studies^{20,22-25)} were not stroke subjects and therefore the results could not be used to apply therapies for stroke management.

From 2005 to 2011, the Committee of Korean standard differentiation of the symptoms and signs for the stroke had been worked in Korea. In

this work, the Committee suggested four types of stroke pattern identification (the Fire Heat Pattern, the Phlegm Dampness Pattern, the Qi Deficiency Pattern, and the Yin Deficiency Pattern) with agreement of experts²⁸⁻³⁰⁾. We conducted binary logistic regression analysis to identify the variables that affected these four pattern-identifications and derive the predictive logistic equations for stroke subjects in 2011 to develop a standardized diagnostic pattern-identification equation for stroke subjects³¹⁾. Where, the predictive logistic equations for the Fire Heat Pattern, the Phlegm Dampness Pattern, the Qi Deficiency Pattern, and the Yin Deficiency Pattern were suggested. However, the sample size of the previous study was relatively small.

In the current study, we aimed to derive new predictive logistic equations for each diagnostic pattern using an expanded number of subjects.

Materials and Methods

1. Subjects

This study was a hospital-based multi-center trial in patients from Korean Medicine university hospitals in the national capital region including Seoul Metropolitan City, and Incheon Metropolitan City as follows: Kyung Hee University Korean Medicine Hospital (Seoul), Kyung Hee University Korean Medicine Hospital at Gangdong (Seoul), and Gil Oriental Medical Hospital of Gacheon University (Incheon). The Institutional Review Board (IRB) approved this study (KOMC IRB 2011-02, KHNMC OHIRB-2012-003, and KWOMH IRB 2006-101).

Subjects with ischemic and hemorrhagic stroke who met International Statistical Classification of Diseases and Related Health Problems 10th

revision code I63 and I61 within 30 days of symptom onset, confirmed by computerized tomography or magnetic resonance imaging were included. Subjects with traumatic stroke such as those with subarachnoid, subdural, and epidural hemorrhage were excluded. Subjects had been enrolled consecutively between April 2007 and December 2012. If there are patients who experienced two or more stroke attack during follow-up period, we only use the data which were obtained in the first admission time. Informed consent was obtained from all study subjects or protectors after a full explanation of the study.

2. Measured Variables

Subjects' general information including medical history including stroke risk factors, the Trial of ORG 10172 in Acute Stroke Treatment (TOAST)^{32,33} classification of the stroke, as well as the variables related to diagnostic pattern-identification (Appendix 1) were measured. All the variables in this study are binary pattern; yes or no. The translation of these medical terms into English was based on the guideline suggested by the World Health Organization (WHO)³⁴.

These subjects' data were collected using Case Report Forms (CRFs) and Standard Operation Procedures (SOPs) made by the expert committee of the Korean Institute of Oriental Medicine (KIOM)^{28-30,35,36} were used. These CRFs and SOPs are the same format used in previous study³¹. These procedures were independently conducted without discussion among KMDs (Korean Medicine Doctors). One KMD recorded CRFs and decided KM diagnostic pattern-identification after examination. Another KMD just decided KM diagnostic pattern-identification

after examination.

3. Determination of Pattern Identification and a New Diagnostic Pattern Identification Equation

The diagnostic pattern of each subject was identified independently by two KMDs, each of whom belonged to one of the study hospital and had 3 years or more clinical experience with stroke management. The categories of identified pattern were the Fire Heat, the Phlegm Dampness, the Qi Deficiency, and the Yin Deficiency Pattern. These categories of identified pattern were suggested by KIOM (Korea Institute of Oriental Medicine)^{28-30,35} and used in previous study³¹. The pattern of each subject was confirmed if the two doctors had the same opinion. Cases where the two doctors had different opinions comprised an unclassified group.

Binary logistic regression was used to detect significant variables which affect the Pattern-Identification decision. To remove the variables that had no statistical significance step-by-step, a backward method was used until a logistic equation composed of absolutely significant variables was reached, and each retained variable considered as an independent factor for the predictive model. Fire Heat, Phlegm Dampness, Qi Deficiency, and Yin Deficiency were studied. The derived logistic equation was as follows: $\log(P/1-P) = \beta_0 + \beta_1 X_1 + \beta_2 X_2 + \dots + \beta_n X_n$. In this equation, X s are independent variables, β s are regression coefficients, and P is the calculated probability of the pattern-identification. According to the β coefficient, there would be positively quantified or negatively quantified variables, i.e., if a subject had more positives, the probability of

Appendix 1. Variables related to Korean Medical Pattern-identification

Overweight: Body Mass Index > 23 (kg/m²)
Insomnia: inability to sleep or abnormal wakefulness
Fatigue: lack of strength
Pale complexion: a white complexion with a hint of blue or gray, often caused by yang collapse or exuberance of cold
Yellow complexion: yellow discoloration of the face, generally suggesting accumulation of dampness
Reddened complexion: a complexion redder than normal, indicating the presence of heat
Darkish complexion: dark discoloration of the face, often occurring in cold syndrome, water retention, or blood stasis
Flushed cheeks: localized flush in the cheeks, indicating yin deficiency
Headache: pain in the head
Eye-congestion: congestion in eyeballs indicating presence of heat
Phlegm rale: an abnormal breathing sound caused by phlegm in the airways
Faint low voice: a voice that is faint and low, scarcely audible
Tongue sore: ulceration in the oral cavity or tongue
Halitosis: bad smell from the mouth
Thirst: feeling of dryness of the mouth with a desire to drink
Night sweating: sweating during sleep that ceases on awakening
Chest discomfort: feeling of stuffiness and fullness in the chest
Nausea: an unpleasant sensation with an urge to vomit
Aversion to heat: strong dislike of heat, also known as heat intolerance
Afternoon tidal fever: fever more marked in the afternoon
Heat in the palms and soles: subjective feverish feeling in the palms and soles
Reversal cold of the extremities: pronounced cold in the extremities up to the knees and elbows, also known as cold extremities
Reddish yellow urine: dark yellow or even reddish urine, indicating heat
Pale tongue: a tongue less red than normal, indicating Qi and blood deficiency
Pale red tongue: a tongue of normal color
Red tongue: a tongue redder than normal, indicating the presence of heat
White fur: a tongue coating that is white in color
Yellow fur: a tongue coating that is yellow in color
Thick fur: a tongue coating where the underlying tongue surface is not visible
Dry fur: a tongue coating that looks dry and feels dry to the touch
Teeth-marked tongue: a tongue with dental indentations on its margin
Enlarged tongue: a tongue that is larger than normal, pale in color and delicate
Mirror tongue: a completely smooth tongue free of coating, like a mirror
Floating pulse: a superficially located pulse that can be felt by light touch and grows faint on hard pressure
Deep pulse: a deeply located pulse that can be felt only when pressing hard
Slow pulse: bradycardia
Rapid pulse: tachycardia
Strong pulse: a general term for strongly beating pulse
Vacuous pulse: a general term for a feeble and void pulse
Thin pulse: a pulse as thin as a silk thread, straight and soft, feeble yet always perceptible upon hard pressure
Slippery pulse: a pulse coming and going smoothly like beads rolling on a plate
Rough pulse: a pulse coming and going, not smoothly, with a small, fine, slow joggling tempo like scraping bamboo with a knife
Flooding pulse: a pulse beating like dashing waves with forceful rising and gradual decline

the pattern-identification would increase, whereas if there were more negatives, the probability of non-pattern-identification would increase. The

statistical analysis was performed using SPSS for Windows, version 16.0 (SPSS Inc., Chicago, Illinois, USA).

Results

1. General Characteristics of Each Pattern Identification

A total of 1,972 stroke patients were recruited. Of these, 721 could not be classified as having a certain pattern because the diagnoses of the two doctors differed. These patients were excluded from the analyses. Among the remaining 1,251 patients, 385 patients (30.8%) had the Fire Heat Pattern, 460 patients (36.8%) the Phlegm Dampness Pattern, 212 patients (16.9%) the Qi Deficiency Pattern, and 194 patients (15.5%) the Yin Deficiency Pattern. The Fire Heat Pattern was more common among men, and the Phlegm Dampness Pattern was most common among patients who exhibited more variables related to

heart disease was exhibited most commonly under the Yin Deficiency Pattern. Most (86.1~94.1%) subjects had ischemic stroke of all subtypes. Among the TOAST subtype classifications^{32,33}, all four diagnostic patterns were similarly represented, with SVO being the most common (Table 1). Herein, only descriptive statistics were applied; comparative statistical analysis was not performed because of the lack of balance in the sample sizes of the pattern groups.

2. Retained Significant Symptoms for Fire Heat Pattern and Identification of Fire Heat Diagnostic Pattern Equation

There were 385 patients (30.8% of the total identified patients) in whom two doctors identified Fire Heat Pattern. Teeth marked tongue,

Table 1. Demographic Data classified by Pattern-identification Subtype

	Fire Heat	Phlegm Dampness	Qi Deficiency	Yin Deficiency
Total number	385	460	212	194
Age, yr	65.6±11.0	66.2±10.9	66.7±12.5	67.7±12.3
Gender, male (%)	277 (71.9)	210 (45.7)	71 (33.5)	98 (50.5)
BMI (kg/m ²)	24.3±2.7	24.7±2.9	22.7±2.9	23.2±4.0
Ischemic stroke	344 (89.4)	433 (94.1)	184 (86.8)	167 (86.1)
TOAST				
LAA (%)	125 (32.5)	106 (23.0)	46 (21.7)	59 (30.4)
CE (%)	25 (6.5)	34 (7.4)	17 (8.0)	19 (9.8)
SVO (%)	186 (48.3)	279 (60.7)	116 (54.7)	83 (42.8)
SOE (%)	8 (2.1)	14 (3.0)	5 (2.4)	6 (3.1)
SUE (%)	0	0	0	0
Medical history				
Hypertension (%)	235 (61.0)	311 (67.6)	134 (63.2)	106 (54.6)
Diabetic mellitus (%)	103 (26.8)	148 (32.2)	53 (25.0)	42 (21.6)
Hyperlipidemia (%)	56 (14.5)	93 (20.2)	35 (16.5)	21 (10.8)
Ischemic heart (%)	24 (6.2)	31 (6.7)	14 (6.6)	18 (9.3)
Previous stroke (%)	93 (24.2)	106 (23.0)	48 (22.6)	51 (26.3)

Values in parentheses refer to %.

BMI: Body Mass Index; TOAST: Trial of Org 10172 in Acute Stroke Treatment; LAA: large artery atherosclerosis; CE: cardioembolism; SVO: small vessel occlusion; SOE: stroke of other determined etiology; SUE: stroke of undetermined etiology

the metabolic syndrome (hypertension, diabetic mellitus, and dyslipidemia). Previous ischemic

vacuous pulse, halitosis, thirst, and heat in the palms and soles showed relatively high

significance in the first analysis, but were removed as the backward method progressed. Finally, overweight, fatigue, yellow complexion, reddened complexion, darkish complexion, tongue sore, aversion to heat, reversal cold of the extremities, reddish yellow urine, pale tongue, yellow fur, thick fur, dry fur, deep pulse, slow pulse, strong pulse, thin pulse, slippery pulse, and flooding pulse all showed highly statistically significant correlations with the Fire Heat pattern and were retained in the equation.

Thus, we derived the predictive logistic equation:
 $\log(P/1-P) = 0.387 \times (\text{Overweight}) - 0.668 \times (\text{Fatigue}) - 0.376 \times (\text{Yellow complexion}) + 1.605 \times (\text{Reddened complexion}) - 0.740 \times (\text{Darkish complexion}) + 0.899 \times (\text{Tongue sore}) + 0.680 \times (\text{Aversion to heat}) - 0.628 \times (\text{Reversal cold of the extremities}) + 0.390 \times (\text{Reddish yellow urine}) - 1.265 \times (\text{Pale tongue}) + 0.383 \times (\text{Yellow fur}) + 0.628 \times (\text{Thick fur}) + 0.378 \times (\text{Dry fur}) - 0.660 \times (\text{Deep pulse}) - 0.507 \times (\text{Slow pulse}) + 0.597 \times (\text{Strong pulse}) - 1.187 \times (\text{Thin pulse}) - 0.762 \times (\text{Slippery pulse}) + 1.099 \times (\text{Flooding pulse}) - 1.280$, where P is the probability of Fire Heat Pattern, and the values of the independent variables would be 0 (absent) or 1 (present). R-square calculated by Nagelkerke's method was 0.405.

3. Retained Significant Symptoms for Phlegm Dampness Pattern and Identification of Phlegm Dampness Diagnostic Pattern Equation

There were 460 patients (36.8% of the total identified patients) in whom two doctors identified Phlegm Dampness Pattern. All of the assessed variables were included in the first logistic regression analysis. Yellowish complexion, reddened complexion, flushed cheeks, nausea,

phlegm rale, halitosis, chest discomfort, reddish yellow urine, red tongue, yellow fur, white fur, deep pulse, vacuous pulse, and aversion to heat exhibited relatively high significance at the first analysis, but were removed as the backward method progressed. Ultimately, overweight, darkish complexion, faint low voice, night sweating, pale tongue, dry fur, slow pulse, rapid pulse, thin pulse, slippery pulse, and flooding pulse all showed high statistical significance and were retained in the equation.

Thus, we derived the predictive logistic equation:
 $\log(P/1-P) = 0.599 \times (\text{overweight}) + 0.850 \times (\text{darkish complexion}) - 0.415 \times (\text{faint low voice}) - 0.476 \times (\text{chest discomfort}) - 0.438 \times (\text{night sweating}) + 0.949 \times (\text{pale tongue}) - 0.767 \times (\text{dry fur}) + 0.668 \times (\text{slow pulse}) + 0.625 \times (\text{rapid pulse}) - 0.550 \times (\text{thin pulse}) + 1.908 \times (\text{slippery pulse}) - 0.920 \times (\text{flooding pulse}) - 2.408$, where P is the probability of Phlegm Dampness Pattern, and the values of the independent variables would be 0 (absent) or 1 (present). R-square calculated by Nagelkerke's method was 0.349.

4. Retained Significant Symptoms for Qi Deficiency Pattern and Identification of Qi Deficiency Diagnostic Pattern Equation

There were 212 patients (16.9% of the total identified patients) in whom two doctors identified Qi Deficiency Pattern. All of the assessed variables were included in the first logistic regression analysis. Headache, red tongue, reddish yellow urine, chest discomfort, heat in the palms and soles, and strong pulse showed relatively high significance at the first analysis, but were removed as the backward method progressed. As a result, overweight, insomnia,

fatigue, pale complexion, reddened complexion, faint low voice, tongue sore, halitosis, thirst, aversion to heat, reversal cold of the extremities, yellow fur, thick fur, dry fur, teeth marked tongue, slow pulse, rapid pulse, vacuous pulse, thin pulse, and slippery pulse all showed high statistical significance and were retained in the equation.

Thus, we derived the predictive logistic equation:

$$\log (P/1-P) = -0.722 \times (\text{overweight}) - 0.663 \times (\text{insomnia}) + 0.662 \times (\text{fatigue}) + 0.923 \times (\text{pale complexion}) - 1.130 \times (\text{reddened complexion}) + 1.089 \times (\text{faint low voice}) - 1.143 \times (\text{tongue sore}) - 0.745 \times (\text{halitosis}) - 0.606 \times (\text{thirst}) - 0.858 \times (\text{aversion to heat}) + 0.546 \times (\text{reversal cold of the extremities}) - 1.026 \times (\text{yellow fur}) - 0.683 \times (\text{thick fur}) - 0.791 \times (\text{dry fur}) + 0.624 \times (\text{teeth marked tongue}) + 0.462 \times (\text{deep pulse}) - 0.659 \times (\text{slow pulse}) - 1.936 \times (\text{rapid pulse}) + 0.906 \times (\text{vacuous pulse}) + 0.979 \times (\text{thin pulse}) - 0.927 \times (\text{slippery pulse}) - 0.791$$
, where the P is the probability of Qi Deficiency Pattern, and the values of the independent variables would be 0 (absent) or 1 (present). R-square calculated by Nagelkerke's method was 0.466.

5. Retained Significant Symptoms for Yin Deficiency Pattern and Identification of Yin Deficiency Diagnostic Pattern Equation

There were 194 patients (15.5% of the total identified patients) in whom two doctors identified Yin Deficiency Pattern. All of the assessed variables were included in the first logistic regression analysis. Fatigue, pale complexion, tongue sore, and red tongue showed relatively high significance at the first analysis, but were removed as the backward method

progressed. As a result, overweight, flushed cheeks, faint low voice, halitosis, afternoon tidal fever, night sweating, pale tongue, yellow fur, dry fur, floating pulse, deep pulse, slow pulse, strong pulse, thin pulse, slippery pulse, and flooding pulse all showed high statistical significance and were retained in the equation.

Thus, we derived the predictive logistic equation:

$$\log (P/1-P) = -0.511 \times (\text{overweight}) + 0.621 \times (\text{flushed cheeks}) + 0.464 \times (\text{faint low voice}) - 0.734 \times (\text{halitosis}) + 0.696 \times (\text{afternoon tidal fever}) + 0.426 \times (\text{night sweating}) - 0.702 \times (\text{pale tongue}) - 0.485 \times (\text{yellow fur}) + 0.920 \times (\text{dry fur}) - 0.537 \times (\text{floating pulse}) - 0.801 \times (\text{deep pulse}) - 0.747 \times (\text{slow pulse}) - 0.615 \times (\text{strong pulse}) + 0.539 \times (\text{thin pulse}) - 0.756 \times (\text{slippery pulse}) - 1.188 \times (\text{flooding pulse}) - 1.003$$
, where the P is the probability of Yin Deficiency Pattern, and the values of the independent variables would be 0 (absent) or 1 (present). R-square calculated by Nagelkerke's method was 0.227.

Discussion

The present study was designed to increase the accuracy and reliability of the Traditional Korean Medical Diagnostic Pattern-identification logistic equation model, which was first reported in 2011³¹⁾.

There are two issues at point. First, the information directly obtained by KMDs such as inspection, tongue diagnosis, and pulse diagnosis play more important roles to make the predictive equation than the subjects' subjective complaints. Based on these data, we could know that KMDs are more concerned with the data obtained by physical examination than the data obtained by medical interview in the process of the pattern

identification decision. Second, overweight might be also an important variable in determining diagnostic pattern. In practice, overweight was included in every type of equation. Therefore, body shape could also be a key point in the process of pattern identification decision.

The characteristics of each diagnostic Pattern-Identification derived from this study are as follows. The excessive patterns such as Fire Heat and the Phlegm Dampness Pattern have positively quantified variables in higher proportion than other patterns. Therefore, determining these two patterns could be an inclusive pattern. On the other hand, the deficiency pattern such as Qi and Yin Deficiency Patterns revealed an exclusive pattern composed of many negatively quantified variables.

In this study, we did not investigate the blood stasis, one of the pattern identification types that has been thought to be the main pattern identification of the stroke. The reason for this is as follows. In version 1 of KIOM's study, which established the categories of pattern identification used in the present study, five pattern identification types were included (including blood stasis). However, blood stasis was found in only 24 (2.8%) out of 859 patients in investigation of version 2³⁷⁾. Different clinical circumstances in Korea and China had been noted for this result. In Korea, a medical classic, *Dongeuibogam* has been set as the basic text. Unlike Chinese literature, *Doneuibogam* didn't record blood stasis as one of the pattern identification of the stroke³⁷⁾. Therefore, blood stasis type was excluded in version 3³⁰⁾, and we did not investigate blood stasis according to version 3.

There are some limitations in the present study as well. Inspection, tongue and pulse diagnosis

were obliged to rely on the KMDs' subjective perception. To overcome this, objective measurements should be developed and suggested. Because this is not a community-based study, our results could not be generalized for Korean people. Finally, in this study, we excluded cases which the decision on pattern identification of two experts was different considering the difficulty of statistical processing. As a result, we could not evaluate the situation which two or more pattern identification types were mixed. Future studies need other statistical processing methods that can reflect the situation in which two or more pattern identification are mixed.

The strengths of this study are as follows. First, new predictive logistic diagnostic pattern equations including more significant variables and based on larger samples are coming up. Second, coefficients in diagnostic patterns as derived by binary logistic regression analysis are suggested. Among study groups of traditional medicine in East Asia, there have been several studies to suggest the significant variables for the determination of diagnostic patterns. In Japan, several standardized questionnaires have been suggested: OKETSU (blood stasis)²⁰⁾, Qi Deficiency, Qi movement Stagnation²⁷⁾, Phlegm Dampness, and Cold-Heat Pattern-identification. In these questionnaires, coefficients for each value were suggested. However, only the OKETSU questionnaire by Terasawa et al²⁰⁾ and the Qi movement Stagnation questionnaire by Oketsu²⁷⁾ clearly stated the methodology and development process. In China, there have been similar attempts at standardization of diagnostic pattern-identification. The standard form of pattern-identification for stroke subjects was suggested in 1994²¹⁾. However, the form of this suggestion is only the

list of symptoms. There was no coefficient that could reveal the importance of each symptom in each pattern. In Korea, similar suggestions exist. Kim et al suggested a questionnaire for the Heat Pattern²² and Park et al suggested a questionnaire for the Phlegm Dampness Pattern²⁵. However, both of them were developed using the Delphi method from KMD clinicians. In 2012, Lee et al established five stroke patterns from the traditional Korean Medical literature and extracted 117 indicators required for diagnosis. These indicators were evaluated by a field test and verified by an expert committee however, they did not suggest coefficients²⁶. In this study, binary logistic regression analysis was applied to derive significant variables for the determination of each diagnostic pattern in order to overcome the limitations of the studies described above.

In this study, standardization of the Traditional Korean Medical Pattern-identification in stroke subjects was intended. After the regression analysis, the predictive logistic equations for Fire Heat, Phlegm Dampness, Qi Deficiency, and Yin Deficiency pattern were determined. These criteria would be useful to determine individual stroke subjects' pattern-identification in the clinical setting. Furthermore, we think the result of this study could be able to be a supportive teaching material for improving Traditional Korean Medicine Doctors' clinical ability for pattern identification.

Aknowledgements

This work was supported by a Grant from the Korean Institute of Oriental Medicine (K09200). This paper is based on Seungwon Kwon's doctoral dissertation.

Competing Interests

The authors declare that they have no competing interests.

References

1. Donnan GA, Fisher M, Macleod M, Davis SM. Stroke. *Lancet*. 2008;371(9624):1612-1623.
2. Wardlaw JM, Murray V, Berge E, del Zoppo G, Sandercock P, Lindley RL, et al. Recombinant tissue plasminogen activator for acute ischaemic stroke: an updated systematic review and meta-analysis. *Lancet*. 2012;379(9834):2364-2372.
3. DeMers G, Meurer WJ, Shih R, Rosenbaum S, Vilke GM. Tissue plasminogen activator and stroke: review of the literature for the clinician. *J Emerg Med*. 2012;43(6):1149-54.
4. ESPS Group. European Stroke Prevention Study. *Stroke*. 1990;21(8):1122-1130.
5. Gong S, Lin C, Zhang D, Kong X, Chen J, Wang C, et al. Effects of Intensive Blood Pressure Reduction on Acute Intracerebral Hemorrhage: A Systematic Review and Meta-analysis. *Neurology. Sci Rep*. 2017 Sep 6;7(1):10694.
6. Hemphill JC 3rd, Greenberg SM, Anderson CS, Becker K, Bendok BR, Cushman M, et al. Guidelines for the Management of Spontaneous Intracerebral Hemorrhage: A Guideline for Healthcare Professionals From the American Heart Association/American Stroke Association. *Stroke*. 2015;46(7):2032 - 2060.
7. Kim JS, Yoon SS. Perspectives of stroke in persons living in Seoul, South Korea: a survey of 1000 subjects. *Stroke*. 1997;28(6):

- 1165-1169.
8. Jung WS, Choi DJ, Cho KH, Lee KS, Moon SK, Kim YS, et al. Safety and efficacy assessment of chungpyesagan-tang for acute ischemic stroke. *Am J Chin Med.* 2003;31(2): 181-190.
 9. Park SU, Jung WS, Moon SK, Ko CN, Cho KH, Kim YS, et al. Chunghyul-dan (Qingxie-dan) improves arterial stiffness in patients with increased baPWV. *Am J Chin Med.* 2006; 34(4):553-563.
 10. Yun SP, Jung WS, Park SU, Moon SK, Ko CN, Cho KH, et al. Anti-hypertensive effect of chunghyul-dan (qingxue-dan) on stroke patients with essential hypertension. *Am J Chin Med.* 2005;33(3):357-364.
 11. Cho K, Noh K, Jung W, Park S, Moon S, Park J, et al. A preliminary study on the inhibitory effect of Chunghyul-dan on stroke recurrence in patients with small vessel disease. *Neurol Res.* 2008;30(6):655-658.
 12. Jung WS, Kwon S, Cho SY, Park SU, Moon SK, Park JM, et al. The Effects of Chunghyul-Dan (A Korean Medicine Herbal Complex) on Cardiovascular and Cerebrovascular Diseases: A Narrative Review. *Evid Based Complement Alternat Med.* 2016;2016: 2601740.
 13. Moon SK, Whang YK, Park SU, Ko CN, Kim YS, Bae HS, et al. Antispastic effect of electroacupuncture and moxibustion in stroke patients. *Am J Chin Med.* 2003;31(3):467-474.
 14. Kim YS, Hong JW, Na BJ, Park SU, Jung WS, Moon SK, et al. The effect of low versus high frequency electrical acupoint stimulation on motor recovery after ischemic stroke by motor evoked potentials study. *Am J Chin Med.* 2008;36(1):45-54.
 15. Goto H, Satoh N, Hayashi Y, Hikiami H, Nagata Y, Obi R, et al. A chinese herbal medicine, tokishakuyakusan, reduces the worsening of impairments and independence after stroke: a 1-year randomized, controlled trial. *Evid Based Complement Alternat Med.* 2011;2011:194046.
 16. Venketasubramanian N, Young SH, Tay SS, Umapathi T, Lao AY, Gan HH, et al. CHinese Medicine NeuroAiD Efficacy on Stroke Recovery - Extension Study (CHIMES-E): A Multicenter Study of Long-Term Efficacy. *Cerebrovasc Dis.* 2015;39(5-6):309-318.
 17. Jung W, Kwon S, Park S, Moon S. Can combination therapy of conventional and oriental medicine improve poststroke aphasia? Comparative, observational, pragmatic study. *Evid Based Complement Alternat Med.* 2012;2012:654604.
 18. Seo YR, Jung WS, Park SU, Moon SK, Park JM, Park JY.. The effect of ouhyul herbal acupuncture point injections on shoulder pain after stroke. *Evid Based Complement Alternat Med.* 2013;2013:504686.
 19. Wu HZ, Fang ZQ, Cheng PJ. Introduction to Diagnosis in Traditional Chinese Medicine. World Century Publishing Corporation, Hackensack, NJ, USA, 2013;69.
 20. Terasawa K, Shinoda H, Imadaya A, Tosa H, Bandoh M, Satoh N. The presentation of diagnostic criteria for Oketsu syndrome. *Japanese. Kampo Med.* 1983;34(1):1-17.
 21. State Administration of Traditional Chinese Medicine of the People's Republic of China. A standard diagnosis for stroke in TCM. *J Beijing University Trad Chin Med.* 1994;17: 64-66.
 22. Kim SK, Park YB. Development of questionnaires

- for cold-heat patternization. *J Korean Inst Oriental Med Diag.* 2003;7(1):64-75.
23. Yang DH, Park YJ, Park YB, Lee SC. Development of questionnaires for blood stasis pattern. *J Korean Inst Oriental Med Diag.* 2006;10(1):141-152.
 24. Bae NS, Yang DH, Park YJ, Park YB, Lee SC, Oh HS. Development of questionnaires for Yol patternization. *J Korean Inst Oriental Med Diag.* 2006(1);10:98-108.
 25. Park JS, Yang DH, Kim MY, Lee SC, Park YJ, Park YB. Development of questionnaire for Damum patternization. *J Korean Inst Oriental Med Diag.* 2006;10(1):64-77.
 26. Lee JA, Park TY, Lee J, Moon TW, Choi J, Kang BK, et al. Developing indicators of pattern identification in patients with stroke using traditional Korean medicine. *BMC Res Notes.* 2012;5:136.
 27. Okitsu R, Iwasaki K, Monma Y, Takayama S, Kaneko S, Shen G, et al. Development of a questionnaire for the diagnosis of Qi stagnation. *Complement Ther Med.* 2012;20(4): 207-217.
 28. Kim JK, Seol IC, Lee I, Jo HK, Yu BC, Choi SM. Report on the Korean standard differentiation of the symptoms and signs for the stroke-1. *Korean J Oriental Physiology & Pathology.* 2006;20(1):229-234.
 29. Go HY, Kim YK, Kang BK, Kim BY, Ko MM, Kang KW, et al. Report on the Korean standard differentiation of the symptoms and signs for the stroke-2. *Korean J Oriental Physiology & Pathology.* 2006;20(6):1789-1791.
 30. Lee JA, Lee JS, Kang BK, Ko MM, Mun TU, Cho KH, et al. Report on the Korean Standard Pattern Identifications for Stroke-III. *Korean J Orient Int Med.* 2011;32(2):232-242.
 31. Kim HJ, Bae HS, Park SU, Moon SK, Park JM, Jung WS. Clinical approach to the standardization of oriental medical diagnostic pattern identification in stroke patients. *Evid Based Complement Alternat Med.* 2011;2011.
 32. Adams HP Jr, Bendixen BH, Kappelle LJ, Biller J, Love BB, Gordon DL, et al. Classification of subtype of acute ischemic stroke. Definitions for use in a multicenter clinical trial. TOAST. Trial of Org 10172 in Acute Stroke Treatment. *Stroke.* 1993;24(1): 35-41.
 33. Gordon DL, Bendixen BH, Adams HP Jr, Clarke W, Kappelle LJ, Woolson RF. Interphysician agreement in the diagnosis of subtypes of acute ischemic stroke: implications for clinical trials. The TOAST Investigators. *Neurology.* 1993;43(5):1021-1027.
 34. WHO International Standard Terminologies on Traditional Medicine in the Western Pacific Region. WHO Western Pacific Regional Office, 2007.
 35. Lee MG, Kang BK, Kim BY, Ko HY, Choi SM, Seol IC, et al. Improvement of tentative Korean standard differentiation of the symptoms and signs for stroke for clinical application. *Korean J Oriental Physiology & Pathology.* 2007;21(1):347-351.
 36. Ko SG, Jun CY, Park JH, Han CH, Ko HY, Yoon Y, et al. Development of CRF and SOPs for the identification of risk factors of the cerebrovascular diseases in the east and west medicine. *J Korean Oriental Med.* 2006;27(1):204-219.
 37. Lee JS, Kim SY, Kang BK, Ko MM, Kim JC, Oh DS, et al. A Review of Static Blood Pattern in Stroke Pattern Diagnosis. *Korean J Orient Int Med.* 2009;30(4):813-820.

ORCID

Woo-Sang Jung	https://orcid.org/0000-0001-7355-7684
Seung-Yeon Cho	https://orcid.org/0000-0003-3149-9759
Seong-Uk Park	https://orcid.org/0000-0002-4617-2719
Sang-Kwan Moon	https://orcid.org/0000-0003-0497-3080
Jung-Mi Park	https://orcid.org/0000-0002-0851-4431
Chang-Nam Ko	https://orcid.org/0000-0002-4749-7961
Ki-Ho Cho	https://orcid.org/0000-0001-5824-9114
Seungwon Kwon	https://orcid.org/0000-0002-1857-3515